

ON THE STRUCTURE OF CATEGORICAL ABSTRACT ELEMENTARY CLASSES WITH AMALGAMATION

MONICA M. VANDIEREN AND SEBASTIEN VASEY

ABSTRACT. For \mathcal{K} an abstract elementary class with amalgamation and no maximal models, we show that categoricity in a high-enough cardinal implies structural properties such as the uniqueness of limit models and the existence of good frames. This improves several classical results of Shelah.

Theorem 0.1. *Let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in a $\lambda \geq \beth_{(2^\mu)^+}$, then:*

- (1) *Whenever $M_0, M_1, M_2 \in \mathcal{K}_\mu$ are such that M_1 and M_2 are limit over M_0 , we have $M_1 \cong_{M_0} M_2$.*
- (2) *If $\mu > \text{LS}(\mathcal{K})$, the model of size λ is μ -saturated.*
- (3) *If $\mu \geq \beth_{(2^{\text{LS}(\mathcal{K})})^+}$ and $\lambda \geq \beth_{(2^\mu)^+}$, then there exists a type-full good μ -frame with underlying class the saturated models in \mathcal{K}_μ .*

Our main tool is the symmetry property of splitting (previously isolated by the first author). The key lemma deduces symmetry from failure of the order property.

CONTENTS

1.	Introduction	2
2.	Background	5
3.	Symmetry from no order property	8
4.	Symmetry and categoricity	13
5.	Good frames and weak tameness	16
	References	18

Date: September 15, 2015

AMS 2010 Subject Classification: Primary 03C48. Secondary: 03C45, 03C52, 03C55, 03C75, 03E55.

Key words and phrases. Abstract elementary classes; Categoricity; Superstability; Tameness; Symmetry; Splitting; Good frame; Limit models; Saturated models.

The second author is supported by the Swiss National Science Foundation.

1. INTRODUCTION

The guiding conjecture for the classification of abstract elementary classes (AECs) is Shelah’s categoricity conjecture. Most progress towards this conjecture has been made under the assumption that the categoricity cardinal is a successor, e.g. [She99, GV06a, Bon14]¹. In this paper, we assume the amalgamation property and no maximal models and deduce new structural results without having to assume that the categoricity cardinal is a successor, or even “high-enough” cofinality.

Consider an AEC \mathcal{K} with amalgamation and no maximal models which is categorical in a cardinal $\lambda > \text{LS}(\mathcal{K})$. Then \mathcal{K} is stable in every cardinal below λ [She99, Claim 1.7.(b)]; so if $\text{cf}(\lambda) > \text{LS}(\mathcal{K})$, then the model of size λ is $\text{cf}(\lambda)$ -saturated². In particular, if λ is regular then the model of size λ is saturated. Baldwin [Bal09, Problem D.1.(2)] has asked if this generalizes to any “sufficiently large” cardinal λ , so let us discuss what happens if we have no control over the cofinality of λ .

One strategy to show saturation of the model of size λ is to show that \mathcal{K} is stable in λ . However an example of Hart and Shelah [HS90] yields (for an arbitrary $k < \omega$) a sentence $\psi_k \in L_{\omega_1, \omega}$ categorical in $\aleph_0, \aleph_1, \dots, \aleph_k$, but not stable in \aleph_k , [BK09] or see [Bal09, Corollary 26.5.4]. Therefore it is not in general true that categoricity in λ implies stability in λ .

On the other hand, it *is* true if we assume a locality property for Galois types: tameness. This is due to the second author and combines the stability transfer in [Vasa] and the Shelah-Villaveces theorem [SV99].

Fact 1.1. *If \mathcal{K} is a $\text{LS}(\mathcal{K})$ -tame AEC with amalgamation, has no maximal models, and is categorical in a $\lambda > \text{LS}(\mathcal{K})$, then \mathcal{K} is stable in every cardinal. In particular, the model of size λ is saturated.*

Proof. By the Shelah-Villaveces theorem (see Fact 4.4), \mathcal{K} is $\text{LS}(\mathcal{K})$ -superstable (see Definition 2.3). Let $\mu \geq \text{LS}(\mathcal{K})$. By [Vasb, Proposition 10.10], \mathcal{K} is μ -superstable, so in particular stable in μ . \square

¹Recently, the second author has proved a categoricity transfer theorem without assuming that the categoricity cardinal is a successor, but assuming that the class is universal [Vasc] (other partial results not assuming categoricity in a successor cardinal are in [Vasb] and [She09]).

²In the sense of Galois types, i.e. M is λ -saturated if every Galois type over a model of size less than λ contained in M is realized in M . This is called “ λ -Galois-saturated” by some authors, but here we always drop the “Galois” (and similarly for other concepts such as stability).

In this paper, we do *not* assume tameness: we show that we can instead take λ sufficiently big (this is (1) of the theorem in the abstract, see Corollary 4.7 for the proof):

Theorem 1.2. *Let \mathcal{K} be an AEC with amalgamation and no maximal models and let $\mu > \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in a $\lambda \geq \beth_{(2^\mu)^+}$, then the model of size λ is μ -saturated.*

Note that we only obtain μ -saturation, not full saturation (the slogan is that categoricity cardinals above $\beth_{(2^\mu)^+}$ behave as if they had cofinality at least μ). However if $\lambda = \beth_\lambda$ this gives full saturation (this is used to obtain a downward categoricity transfer, see Corollary 4.9). Moreover μ -saturation is enough for many applications, as many of the results of [She99] only assume categoricity in a λ with $\text{cf}(\lambda) > \mu$ (for a fixed $\mu \geq \text{LS}(\mathcal{K})$). For example, we show how to obtain *weak tameness* (i.e. tameness over saturated models) from categoricity in a big-enough cardinal (this is Theorem 4.14). We can then build a local notion of independence: a good μ -frame (this is (3) of the theorem in the abstract, see Corollary 5.4 for a proof):

Theorem 1.3. *Let $\mu \geq \beth_{(2^{\text{LS}(\mathcal{K})})^+}$. Assume that \mathcal{K} is categorical in some $\lambda \geq \beth_{(2^{\mu^+})^+}$. Then there exists a type-full good μ -frame with underlying class the saturated models in \mathcal{K}_μ .*

This improves on [Vasa, Theorem 7.4], which assumed categoricity in a successor (and a higher Hanf number bound). This also (partially) answers [She99, Remark 4.9.(1)] which asked whether there is a parallel to forking in categorical AECs with amalgamation.

The key to the proof of Theorem 1.2 is a close study of the symmetry property for splitting, identified by the first author in [Vana]. There it was shown (assuming superstability in μ) that symmetry of μ -splitting is equivalent to the continuity of reduced towers of size μ , which itself implies uniqueness of limit models in μ . It was also shown that symmetry of μ -splitting follows from categoricity in μ^+ . In [VV], we improved this by only requiring categoricity in a λ of cofinality bigger than μ :

Fact 1.4 (Corollary 5.2 in [VV]). *Let \mathcal{K} be an AEC with amalgamation and no maximal models. Let $\mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is categorical in a cardinal λ with $\text{cf}(\lambda) > \mu$. Then \mathcal{K} is μ -superstable and has μ -symmetry. In particular [VV, Theorem 0.1], it has uniqueness of limit models in μ : for any $M_0, M_1, M_2 \in \mathcal{K}_\mu$, if M_1 and M_2 are limit over M_0 , then $M_1 \cong_{M_0} M_2$.*

Here we replace the cofinality assumption on the categoricity cardinal with the assumption that the categoricity cardinal is big enough (this also proves (2) of the theorem in the abstract, see Theorem 4.5):

Theorem 1.5. *Let \mathcal{K} be an AEC with amalgamation and no maximal models. Let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in a $\lambda \geq \beth_{(2^\mu)^+}$, then \mathcal{K} is μ -superstable and has μ -symmetry. In particular, \mathcal{K} has uniqueness of limit models in μ .*

Remark 1.6. *This gives a proof (assuming amalgamation and a high-enough categoricity cardinal) of the (in)famous [SV99, Theorem 3.3.7], where a gap was identified in the first author's Ph.D. thesis. The gap was fixed assuming categoricity in μ^+ in [Van06, Van13] (see also the exposition in [GVV]). In [BG, Corollary 6.10], this was improved to categoricity in an arbitrary $\lambda > \mu$ provided that μ is big-enough and the class satisfies strong locality assumptions (full tameness and shortness and the extension property for coheir). In [Vasa, Theorem 7.11], only tameness was required but the categoricity had to be in a λ with $\text{cf}(\lambda) > \mu$. Still assuming tameness, this is shown for categoricity in any $\lambda \geq \beth_{(2^\mu)^+}$ in [BV, Theorem 7.1] and in any $\lambda > \mu$ in [VV, Theorem 6.4]. Finally without tameness the result was proved for any λ with $\text{cf}(\lambda) > \mu$ in [VV, Theorem 0.1].*

The proof of Theorem 1.5 is technical but conceptually not hard: we show that a failure of μ -symmetry would give the order property, which in turn would imply instability below the categoricity cardinal. The idea of using the order property to prove symmetry of an independence relation is due to Shelah, [She75, Theorem 6.10(ii)] or see [She90, Theorem III.4.13]. In [BGKV], an abstract generalization of Shelah's proof to any independence notion satisfying extension and uniqueness was given. Here we adapt the proof of [BGKV] to splitting. This uses the extension property of splitting for models of different sizes from [VV].

In general, we obtain that an AEC with amalgamation categorical in a high-enough cardinal has many structural properties that were previously only known for AECs categorical in a cardinal of high-enough *cofinality*, or even just in a successor. This improves several classical results from Shelah's milestone study of categorical AECs with amalgamation [She99].

This paper was written while the second author was working on a Ph.D. thesis under the direction of Rami Grossberg at Carnegie Mellon University and he would like to thank Professor Grossberg for his guidance and assistance in his research in general and in this work specifically.

2. BACKGROUND

Throughout this paper, we assume:

Hypothesis 2.1. \mathcal{K} is an AEC with amalgamation.

For convenience, we fix a big-enough monster model \mathfrak{C} and work inside \mathfrak{C} . This is possible since by Remark 2.4, we will have the joint embedding property in addition to the amalgamation property for models of the relevant cardinalities.

Many of the pre-requisite definitions and notations used in this paper can be found in [GVV]. Here we recall the more specialized concepts that we use explicitly. We begin by recalling the definition of non-splitting, a notion of independence from [She99, Definition 3.2].

Definition 2.2. A type $p \in \text{ga-S}(N)$ does not μ -split over M if and only if for any $N_1, N_2 \in \mathcal{K}_\mu$ such that $M \leq N_\ell \leq N$ for $\ell = 1, 2$, and any $f : N_1 \cong_M N_2$, we have $f(p \upharpoonright N_1) = p \upharpoonright N_2$

The definition of superstability below is already implicit in [SV99] and has since then been studied in several papers, e.g. [Van06, GVV, Vasb, BV, GV, VV]. We will use the formulation from [Vasb, Definition 10.1]:

Definition 2.3. \mathcal{K} is μ -superstable (or superstable in μ) if:

- (1) $\mu \geq \text{LS}(\mathcal{K})$.
- (2) \mathcal{K}_μ is nonempty, has joint embedding, and no maximal models.
- (3) \mathcal{K} is stable in μ^3 , and:
- (4) μ -splitting in \mathcal{K} satisfies the following locality (sometimes called continuity) and “no long splitting chains” properties:

For any limit ordinal $\alpha < \mu^+$, for every sequence $\langle M_i \mid i < \alpha \rangle$ of models of cardinality μ with M_{i+1} universal over M_i and for every $p \in \text{ga-S}(\bigcup_{i < \alpha} M_i)$, we have that:

- (a) If for every $i < \alpha$, the type $p \upharpoonright M_i$ does not μ -split over M_0 , then p does not μ -split over M_0 .
- (b) There exists $i < \alpha$ such that p does not μ -split over M_i .

Remark 2.4. By our global hypothesis of amalgamation (Hypothesis 2.1), if \mathcal{K} is μ -superstable, then $\mathcal{K}_{\geq \mu}$ has joint embedding.

Remark 2.5. By the weak transitivity property of μ -splitting [Vas, Proposition 3.7], condition (4a) could have been omitted (i.e. it follows from the rest).

³That is, $|\text{ga-S}(M)| \leq \mu$ for all $M \in \mathcal{K}_\mu$. Some authors call this “Galois-stable”.

The main tool of this paper is the concept of symmetry over limit models which was identified in [Vana]:

Definition 2.6. *An abstract elementary class exhibits symmetry for μ -splitting (or μ -symmetry for short) if whenever models $M, M_0, N \in \mathcal{K}_\mu$ and elements a and b satisfy the conditions 1-4 below, then there exists M^b a limit model over M_0 , containing b , so that $\text{ga-tp}(a/M^b)$ does not μ -split over N . See Figure 1.*

- (1) M is universal over M_0 and M_0 is a limit model over N .
- (2) $a \in M \setminus M_0$.
- (3) $\text{ga-tp}(a/M_0)$ is non-algebraic and does not μ -split over N .
- (4) $\text{ga-tp}(b/M)$ is non-algebraic and does not μ -split over M_0 .




FIGURE 1. A diagram of the models and elements in the definition of symmetry. We assume the type $\text{ga-tp}(b/M)$ does not μ -split over M_0 and $\text{ga-tp}(a/M_0)$ does not μ -split over N . Symmetry implies the existence of M^b a limit model over M_0 containing b , so that $\text{ga-tp}(a/M^b)$ does not μ -split over N .

We recall a few results of the first author showing the importance of the symmetry property:

Fact 2.7 (Theorem 5 in [Vana]). *If \mathcal{K} is μ -superstable and has μ -symmetry, then for any $M_0, M_1, M_2 \in \mathcal{K}_\mu$, if M_1 and M_2 are limit models over M_0 , then $M_1 \cong_{M_0} M_2$.*

For $\lambda > \text{LS}(\mathcal{K})$, we will write $\mathcal{K}^{\lambda\text{-sat}}$ for the class of λ -saturated models in $\mathcal{K}_{\geq \lambda}$ (we order it with the strong substructure relation induced by \mathcal{K}). By [Vanb] superstability and symmetry together imply that the union of certain chains of saturated models is saturated. This has an easier formulation in [VV, Theorem 6.6]:

Fact 2.8. *Assume \mathcal{K} is μ -superstable, μ^+ -superstable, and has μ^+ -symmetry. Then $\mathcal{K}^{\mu^+ \text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\mu^+ \text{-sat}}) = \mu^+$.*

The following downward transfer of the symmetry property was the key result of [VV]:

Fact 2.9 (Theorem 1.1 in [VV]). *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, \lambda]$. If \mathcal{K} has λ -symmetry, then \mathcal{K} has μ -symmetry.*

It will be convenient to use the following independence notion. A minor variation (where “limit over” is replaced by “universal over”) appears in [Vasa, Definition 3.8].

Definition 2.10 (Definition 3.1 in [VV]). *Let $M_0 \leq M \leq N$ be models with $M_0 \in \mathcal{K}_\mu$. We say a type $p \in \text{ga-S}(N)$ explicitly does not μ -fork over (M_0, M) if:*

- (1) *M is limit over a model containing M_0 .*
- (2) *p does not μ -split over M_0 .*

We say that p does not μ -fork over M if there exists M_0 so that p explicitly does not μ -fork over (M_0, M) .

Remark 2.11. *Assuming μ -superstability, the relation “ p does not μ -fork over M ” is very close to defining an independence notion with the properties of forking in a first-order superstable theory (i.e. a good μ -frame, see Section 5). In fact using tameness (or just, as we will show in Section 5, weak tameness) it can be used to do precisely that, see [Vasa].*

μ -Forking has the following properties:

Fact 2.12. *Let $\mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is stable in μ .*

- (1) [Van06, I.4.10, I.4.12] *Extension and uniqueness in μ : Let $M \leq N$ be in \mathcal{K}_μ , and $p \in \text{ga-S}(M)$ explicitly does not μ -fork over (M_0, M) , then there exists a unique $q \in \text{ga-S}(N)$ extending p so that q explicitly does not μ -fork over (M_0, M) . Moreover q is algebraic if and only if p is.*
- (2) [VV, Proposition 4.3] *Continuity above μ : Assume that \mathcal{K} is μ -superstable. Suppose $\langle M_i \in \mathcal{K}_{\geq \mu} \mid i < \delta \rangle$ is an increasing sequence of models so that, for all $i < \delta$, M_{i+1} is universal over M_i . Let $p \in \text{ga-S}(\bigcup_{i < \delta} M_i)$. If $p \upharpoonright M_i$ does not μ -split over M_0 for each $i < \delta$, then p does not μ -split over M_0 .*
- (3) *Extension above μ : Let $\lambda > \mu$. Let $M \leq N$ be in $\mathcal{K}_{[\mu, \lambda]}$. Let $p \in \text{ga-S}(M)$ be such that p explicitly does not μ -fork over (M_0, M) .*

If \mathcal{K} is superstable in every $\chi \in [\mu, \lambda]$, then there exists $q \in \text{ga-S}(N)$ extending p and explicitly not μ -forking over (M_0, M) . Moreover q is algebraic if and only if p is.

Proof of (3). By induction on $\|N\|$. Let a realize p . If $\|N\| = \|M\|$ this is given by [VV, Proposition 4.4]. If $\|M\| < \|N\|$, build $\langle N_i \in \mathcal{K}_{\|M\|+|i|} : i \leq \|N\| \rangle$ increasing continuous such that $N_0 = M$, N_{i+1} is limit over N_i , and $\text{ga-tp}(a/N_i)$ explicitly does not μ -fork over (M_0, M) . This is possible by the induction hypothesis and the continuity property of forking. Now N_λ is $\|N\|$ -universal over $N_0 = M$, so let $f : N \xrightarrow{M} N_\lambda$. Let $q := f^{-1}(\text{ga-tp}(a/f[N]))$. It is easy to check that q is as desired. \square

3. SYMMETRY FROM NO ORDER PROPERTY

In this section we show (assuming enough instances of superstability) that the negation of symmetry implies the order property, and hence contradicts stability. This is similar to [BGKV, Theorem 5.14], but due to the intricate definition of the symmetry property for splitting, some technical details have to be handled. We first give an equivalent definition of symmetry. Recall that in [VV, Definition 3.3], we gave three variations on the symmetry property:

- (1) The uniform μ -symmetry, which is essentially Definition 2.6 (and in fact is formally equivalent to it).
- (2) The non-uniform μ -symmetry, which weakens the conclusion of uniform μ -symmetry by “changing” the model N that $\text{ga-tp}(a/M^b)$ does not μ -split over.
- (3) The weak non-uniform μ -symmetry which strengthens the hypotheses of non-uniform μ -symmetry by requiring that $\text{ga-tp}(b/M)$ does not μ -fork, instead of μ -split over M_0 .

There is a fourth possible variation, the weak uniform μ -symmetry property, which strengthens the hypotheses of uniform μ -symmetry similarly to the weak non-uniform μ -symmetry, but leaves the conclusion unchanged. For clarity we have underlined the differences between the weak and non-weak definitions.

Definition 3.1. *Let $\mu \geq \text{LS}(\mathcal{K})$.*

- (1) \mathcal{K} has uniform μ -symmetry if for any limit models N, M_0, M in \mathcal{K}_μ where M is limit over M_0 and M_0 is limit over N , if $\text{ga-tp}(b/M)$ does not μ -split over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ explicitly does not μ -fork over (N, M_0) , there exists $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ explicitly does not μ -fork over (N, M_0) .

- (2) \mathcal{K} has weak uniform μ -symmetry if for any limit models N, M_0, M in \mathcal{K}_μ where M is limit over M_0 and M_0 is limit over N , if $\text{ga-tp}(b/M)$ does not μ -fork over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ explicitly does not μ -fork over (N, M_0) , there exists $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ explicitly does not μ -fork over (N, M_0) . See Figure 2.

As noted above, the difference is that in the non-weak version, $\text{ga-tp}(b/M)$ does not μ -split only over M_0 , but in the weak version we require that $\text{ga-tp}(b/M)$ does not μ -fork over M_0 . Thus in the weak version, there exists M'_0 such that M_0 is limit over M'_0 and $\text{ga-tp}(b/M)$ does not μ -split over M'_0 (but we do not ask that M'_0 has any relationship to N).




FIGURE 2. A diagram of the models and elements in the definition of weak uniform μ -symmetry. We require that $\text{ga-tp}(b/M)$ does not μ -fork over M_0 in the weak version, so there exists M'_0 such that M_0 is limit over M'_0 and $\text{ga-tp}(b/M)$ does not μ -split over M'_0

We start by showing that assuming μ -superstability this distinction is inessential, i.e. the two properties are equivalent. We will use the following characterization of symmetry:

Fact 3.2 (Theorem 5 in [Vana]). *Assume that \mathcal{K} is μ -superstable. The following are equivalent:*

- (1) \mathcal{K} has μ -symmetry.
- (2) Reduced towers of size μ are continuous⁴.

⁴See [GVV] for the definitions of these terms.

Lemma 3.3. *Assume that \mathcal{K} is μ -superstable. The weak uniform μ -symmetry is equivalent to uniform μ -symmetry which is equivalent to μ -symmetry.*

Proof. That uniform μ -symmetry is equivalent to μ -symmetry is easy (it appears as [VV, Proposition 3.5]). Clearly, uniform implies weak uniform. Now assuming weak uniform symmetry, the proof of (1) \Rightarrow (2) of Fact 3.2 still goes through. The point is that whenever we consider $\text{ga-tp}(b/M)$ in the proof, $M = \bigcup_{i < \delta} M_i$ for some increasing continuous $\langle M_i : i < \delta \rangle$ with M_{i+1} universal over M_i for all $i < \delta$, and we simply use that by superstability $\text{ga-tp}(b/M)$ does not μ -split over M_i for some $i < \delta$. However we also have that $\text{ga-tp}(b/M)$ explicitly does not μ -fork over (M_i, M_{i+1}) .

Therefore reduced towers are continuous, and hence by Fact 3.2 \mathcal{K} has μ -symmetry. \square

Next we recall the definition of the order property in AECs [She99, Definition 4.3].

Definition 3.4. *Let α and λ be cardinals. A model $M \in \mathcal{K}$ has the α -order property of length λ if there exists $\langle \bar{a}_i : i < \lambda \rangle$ inside M with $\ell(\bar{a}_i) = \alpha$ for all $i < \lambda$, such that for any $i_0 < j_0 < \lambda$ and $i_1 < j_1 < \lambda$, $\text{ga-tp}(\bar{a}_{i_0}\bar{a}_{j_0}/\emptyset) \neq \text{ga-tp}(\bar{a}_{j_1}\bar{a}_{i_1}/\emptyset)$.*

We say that \mathcal{K} has the α -order property of length λ if some $M \in \mathcal{K}$ has it. We say that \mathcal{K} has the α -order property if it has the α -order property of length λ for all cardinals λ .

We will use two important facts: the first says that it is enough to look at length up to the Hanf number. The second that the order property implies instability.

Definition 3.5. *For λ an infinite cardinal, $h(\lambda) := \beth_{(2^\lambda)^+}$.*

Fact 3.6 (Claim 4.5.3 in [She99]). *Let α be a cardinal. If \mathcal{K} has the α -order property of length λ for all $\lambda < h(\alpha + \text{LS}(K))$, then \mathcal{K} has the α -order property.*

Fact 3.7. *If \mathcal{K} has the α -order property and $\mu \geq \text{LS}(\mathcal{K})$ is such that $\mu = \mu^\alpha$, then \mathcal{K} is not stable in μ .*

Proof. By [She99, Claim 4.8.2] (see [BGKV, Fact 5.13] for a proof), there exists $M \in \mathcal{K}_\mu$ such that $|\text{ga-S}^\alpha(M)| > \mu$. By [Bon, Theorem 3.1], \mathcal{K} is not stable in μ . \square

The following lemma appears in some more abstract form in [BGKV, Lemma 5.6]. The lemma says that if we assume that p does not μ -fork over M , then in the definition of non-splitting (Definition 2.2) we can

replace the N_ℓ by arbitrary sequences in N of length at most μ . In the proof of Lemma 3.9, this will be used for sequences of length one.

Lemma 3.8. *Let $\mu \geq \text{LS}(\mathcal{K})$. Let $M \in \mathcal{K}_\mu$ and $N \in \mathcal{K}_{\geq \mu}$ be such that $M \leq N$. Assume that \mathcal{K} is stable in μ . If $p \in \text{ga-S}(N)$ does not μ -fork over M (Definition 2.10), a realizes p , and $\bar{b}_1, \bar{b}_2 \in {}^{\leq \mu}N$ are such that $\text{ga-tp}(\bar{b}_1/M) = \text{ga-tp}(\bar{b}_2/M)$, then $\text{ga-tp}(a\bar{b}_1/M) = \text{ga-tp}(a\bar{b}_2/M)$.*

Proof. Pick $N_0 \in \mathcal{K}_\mu$ containing $\bar{b}_1\bar{b}_2$ with $M \leq N_0 \leq N$. Then $p \upharpoonright N_0$ does not μ -fork over M . Replacing N by N_0 if necessary, we can assume without loss of generality that $N \in \mathcal{K}_\mu$. By definition of μ -forking, there exists $M_0 \in \mathcal{K}_\mu$ such that $M_0 \leq M$ and p does not μ -split over M_0 . By the extension and uniqueness property for μ -splitting there exists N' extending N of cardinality μ so that N' is universal over both N and M , and $\text{ga-tp}(a/N')$ does not μ -split over M_0 . Since $\text{ga-tp}(\bar{b}_1/M) = \text{ga-tp}(\bar{b}_2/M)$ and since N' is universal over N , we can find $f : N \xrightarrow[M]{\rightarrow} N'$ so that $f(\bar{b}_1) = \bar{b}_2$. Since $\text{ga-tp}(a/N')$ does not μ -split over M_0 we know $\text{ga-tp}(f(a)/f(N)) = \text{ga-tp}(a/f(N))$. By our choice of f this implies that there exists $g \in \text{Aut}_{f(N)}(\mathfrak{C})$ so that $g(f(a)) = a$, $g \upharpoonright M = \text{id}_M$, and $g(\bar{b}_2) = \bar{b}_1$. In other words $\text{ga-tp}(f(a)\bar{b}_2/M) = \text{ga-tp}(a\bar{b}_1/M)$. Moreover f^{-1} witnesses that $\text{ga-tp}(a\bar{b}_1/M) = \text{ga-tp}(f(a)\bar{b}_2/M)$, which we have seen is equal to $\text{ga-tp}(a\bar{b}_2/M)$. \square

The next lemma shows that failure of symmetry implies the order property. The proof is similar to that of [BGKV, Theorem 5.14], the difference is that we use Lemma 3.8 and the equivalence between symmetry and weak uniform symmetry (Lemma 3.3).

Lemma 3.9. *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, \lambda)$. If \mathcal{K} does not have μ -symmetry, then it has the μ -order property of length λ .*

Proof. By Lemma 3.3, \mathcal{K} does not have weak uniform μ -symmetry. We first pick witnesses to that fact. Pick limit models $N, M_0, M \in \mathcal{K}_\mu$ such that M is limit over M_0 and M_0 is limit over N . Pick b such that $\text{ga-tp}(b/M)$ does not μ -fork over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ explicitly does not μ -fork over (N, M_0) , and there does *not* exist $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ explicitly does not μ -fork over (N, M_0) . We will show that \mathfrak{C} has the μ -order property of length λ .

We build increasing continuous $\langle N_\alpha : \alpha < \lambda \rangle$ and $\langle a_\alpha, b_\alpha, N'_\alpha : \alpha < \lambda \rangle$ by induction so that for all $\alpha < \lambda$:

- (1) $N_\alpha, N'_\alpha \in \mathcal{K}_{\mu+|\alpha|}$.
- (2) N_0 is limit over M and $b \in |N_0|$.

- (3) $\text{ga-tp}(a_\alpha/M_0) = \text{ga-tp}(a/M_0)$ and $a_\alpha \in |N'_\alpha|$.
- (4) $\text{ga-tp}(b_\alpha/M) = \text{ga-tp}(b/M)$ and $b_\alpha \in |N_{\alpha+1}|$.
- (5) N'_α is limit over N_α and $N_{\alpha+1}$ is limit over N'_α .
- (6) $\text{ga-tp}(a_\alpha/N_\alpha)$ explicitly does not μ -fork over (N, M_0) and $\text{ga-tp}(b_\alpha/N'_\alpha)$ does not μ -fork over M_0 .

This is possible. Let N_0 be any model in \mathcal{K}_μ containing M and a and limit over M . At α limits, let $N_\alpha := \bigcup_{\beta < \alpha} N_\beta$. Now assume inductively that N_β has been defined for $\beta \leq \alpha$, and $a_\beta, b_\beta, N'_\beta$ have been defined for $\beta < \alpha$. By extension for splitting, find $q \in \text{ga-S}(N_\alpha)$ that explicitly does not μ -fork over (N, M_0) and extends $\text{ga-tp}(a/M_0)$. Let a_α realize q and pick N'_α limit over N_α containing a_α . Now by extension again, find $q' \in \text{ga-S}(N'_\alpha)$ that does not μ -fork over M_0 and extends $\text{ga-tp}(b/M)$. Let b_α realize q' and pick $N_{\alpha+1}$ limit over N'_α containing b_α .

This is enough. We show that for $\alpha, \beta < \lambda$:

- (1) $\text{ga-tp}(a_\alpha b/M_0) \neq \text{ga-tp}(ab/M_0)$
- (2) If $\beta < \alpha$, $\text{ga-tp}(ab/M_0) \neq \text{ga-tp}(a_\alpha b_\beta/M_0)$.
- (3) If $\beta \geq \alpha$, $\text{ga-tp}(ab/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$.

For (1), observe that $b \in |N_0| \subseteq |N_\alpha|$ and $\text{ga-tp}(a_\alpha/N_\alpha)$ explicitly does not μ -fork over (N, M_0) . Therefore by monotonicity N_α witnesses that there exists $N_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a_\alpha/M_b)$ explicitly does not μ -fork over (N, M_0) . By failure of symmetry and invariance, we must have that $\text{ga-tp}(a_\alpha b/M_0) \neq \text{ga-tp}(ab/M_0)$.

For (2), suppose $\beta < \alpha$. We know that $\text{ga-tp}(a_\alpha/N_\alpha)$ does not μ -fork over M_0 . Since $\beta < \alpha$, $b, b_\beta \in |N_\alpha|$ and $\text{ga-tp}(b/M) = \text{ga-tp}(b_\beta/M)$, we must have by Lemma 3.8 that $\text{ga-tp}(a_\alpha b/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$ ⁵.

Together with (1), this implies $\text{ga-tp}(ab/M_0) \neq \text{ga-tp}(a_\alpha b_\beta/M_0)$.

To see (3), suppose $\beta \geq \alpha$ and recall that (by (1)) $\text{ga-tp}(ab/M_0) = \text{ga-tp}(a_\beta b/M_0)$. We also have that $\text{ga-tp}(b_\beta/N'_\beta)$ does not μ -fork over M_0 . Moreover $\text{ga-tp}(a/M_0) = \text{ga-tp}(a_\alpha/M_0)$, and $a, a_\alpha \in N'_\beta$. By Lemma 3.8 again, $\text{ga-tp}(ab_\beta/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$. This gives us that $\text{ga-tp}(ab/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$.

Now let \bar{d} be an enumeration of M_0 and for $\alpha < \lambda$, let $\bar{c}_\alpha := a_\alpha b_\alpha \bar{d}$. Then (2) and (3) together tell us that the sequence $\langle \bar{c}_\alpha \mid \alpha < \lambda \rangle$ witnesses the μ -order property of length λ . \square

⁵This is really where we use the equivalence between uniform μ -symmetry and weak uniform μ -symmetry: if we only had failure of uniform μ -symmetry, then we would only know that $\text{ga-tp}(b/M)$ does not μ -split over M_0 , so would be unable to use Lemma 3.8.

Theorem 3.10. *Let $\mu \geq \text{LS}(\mathcal{K})$. Then there exists $\lambda < h(\mu)$ such that if \mathcal{K} is superstable in every $\chi \in [\mu, \lambda)$, then \mathcal{K} has μ -symmetry.*

Proof. If \mathcal{K} is unstable in 2^μ , then we can set $\lambda := (2^\mu)^+$ and get a vacuously true statement; so assume that \mathcal{K} is stable in 2^μ . By Fact 3.7, \mathcal{K} does not have the μ -order property. By Fact 3.6, there exists $\lambda < h(\mu)$ such that \mathcal{K} does not have the μ -order property of length λ . By Lemma 3.9, it is as desired. \square

Remark 3.11. *How can one obtain many instances of superstability as in the hypothesis of Theorem 3.10? One way is categoricity, see the next section. Another way is to start with one instance of superstability and transfer it up using tameness. Indeed by [Vasb, Proposition 10.10], if \mathcal{K} is μ -superstable and μ -tame, then it is superstable in every $\mu' \geq \mu$. Thus Theorem 3.10 generalizes [VV, Theorem 6.4] which obtained μ -symmetry from μ -superstability and μ -tameness.*

4. SYMMETRY AND CATEGORICITY

In this section, we apply Theorem 3.10 to categorical classes. We will use notation from Chapter 14 of [Bal09] (recall from Definition 3.5 that $h(\lambda) := \beth_{(2^\lambda)^+}$):

Definition 4.1. *For $n < \omega$, define inductively:*

- (1) $H_0 := \text{LS}(\mathcal{K})$.
- (2) $H_{n+1} := h(H_n)$.

Throughout this section, we assume (in addition to amalgamation) that \mathcal{K} has no maximal models. This is not a big deal because we can always take a tail of the AEC to obtain it:

Fact 4.2 (Proposition 10.13 in [Vasb]). *If \mathcal{K} is an AEC with amalgamation categorical in a $\lambda \geq H_1$, then there exists $\chi < H_1$ such that $\mathcal{K}_{\geq \chi}$ has no maximal models.*

Hypothesis 4.3. *\mathcal{K} is an AEC with amalgamation and no maximal models.*

The following powerful fact has its roots in [SV99, Theorem 2.2.1], where it was proven assuming the generalized continuum hypothesis instead of amalgamation. This is the main tool to obtain superstability from categoricity. Its proof relies on Ehrenfeucht-Mostowski models, which is why we assumed no maximal models.

Fact 4.4 (The Shelah-Villaveces theorem, Theorem 6.3 in [GV]). *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in λ , then \mathcal{K} is μ -superstable.*

Combining this fact with Theorem 3.10, we obtain symmetry from categoricity in a high-enough cardinal:

Theorem 4.5. *Let $\mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is categorical in a $\lambda \geq h(\mu)$. Then \mathcal{K} is μ -superstable and has μ -symmetry.*

Proof. By the Shelah-Villaveces theorem (Fact 4.4), \mathcal{K} is μ' -superstable in every $\mu' \in [\mu, \lambda)$. By Theorem 3.10, \mathcal{K} has μ -symmetry. \square

Combining Theorem 4.5 with consequences of symmetry, we deduce:

Corollary 4.6. *Let $\mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is categorical in a $\lambda \geq h(\mu)$. For any $M_0, M_1, M_2 \in \mathcal{K}_\mu$, if M_1 and M_2 are both limit over M_0 , then $M_1 \cong_{M_0} M_2$.*

Proof. By Theorem 4.5 and Fact 2.7. \square

Corollary 4.7. *Let $\mu > \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is categorical in a $\lambda \geq \sup_{\mu_0 < \mu} h(\mu_0^+)$. Then the model of size λ is μ -saturated.*

Proof. We check that the model of size λ is μ_0^+ -saturated for every $\mu_0 \in [\text{LS}(\mathcal{K}), \mu)$. Fix such a μ_0 . By Theorem 4.5, \mathcal{K} is μ_0 -superstable, μ_0^+ -superstable, and has μ_0^+ -symmetry. By Fact 2.8 $\mathcal{K}^{\mu_0^+ \text{-sat}}$, the class of μ_0^+ -saturated models in $\mathcal{K}_{\geq \mu_0^+}$, is an AEC with Löwenheim-Skolem number μ_0^+ . Since it has arbitrarily large models, it must have a model of size λ , which is unique by categoricity. \square

We immediately obtain a downward categoricity transfer. We will use:

Fact 4.8 (The AEC omitting type theorem, II.1.10 in [She99]). *Let $\lambda > \chi \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in λ and the model of size λ is χ^+ -saturated, then every model in $\mathcal{K}_{\geq h(\chi)}$ is χ^+ -saturated.*

Corollary 4.9. *Let $\mu = \beth_\mu > \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is categorical in a $\lambda > \mu$. Then \mathcal{K} is categorical in μ .*

Proof. By Corollary 4.7, the model of size λ is μ -saturated. By Fact 4.8, every model in $\mathcal{K}_{\geq \mu}$ is μ -saturated. In particular, every model of size μ is saturated. By uniqueness of saturated models, \mathcal{K} is categorical in μ . \square

Remark 4.10. *For Corollary 4.9, Fact 4.2 shows that the no maximal models hypothesis is not necessary.*

Remark 4.11. *Corollary 4.9 appears as [Vasb, Theorem 10.16] with an additional assumption of $(< \mu)$ -tameness.*

We can improve on Corollary 4.9 using the more powerful downward transfer of [She99]. A key concept in the proof is the following variation on tameness (an important locality for Galois types isolated by Grossberg and the first author in [GV06b]). We use the notation in [Bal09, Definition 11.6].

Definition 4.12. *Let χ, μ be cardinals with $\text{LS}(\mathcal{K}) \leq \chi \leq \mu$. \mathcal{K} is (χ, μ) -weakly tame if for any saturated $M \in \mathcal{K}_\mu$, any $p, q \in \text{ga-S}(M)$, if $p \neq q$, there exists $M_0 \in \mathcal{K}_\chi$ with $M_0 \leq M$ and $p \upharpoonright M_0 \neq q \upharpoonright M_0$.*

The importance of weak tameness is that it is known to follow from categoricity in a suitable cardinal: this appears as [She99, Main Claim II.2.3] and a simplified improved argument is in [Bal09, Theorem 11.15].

Fact 4.13. *Let $\lambda > \mu \geq H_1$. Assume that \mathcal{K} is categorical in λ , and the model of cardinality λ is μ^+ -saturated. Then there exists $\chi < H_1$ such that \mathcal{K} is (χ, μ) -weakly tame.*

We obtain:

Theorem 4.14. *Let $\mu \geq \text{LS}(\mathcal{K})$. Let $\lambda \geq h(\mu^+)$. If \mathcal{K} is categorical in λ , then there exists $\chi < H_1$ such that \mathcal{K} is (χ, μ) -weakly tame.*

Proof. By Corollary 4.7, the model of size λ is μ^+ -saturated. Now apply Fact 4.13. □

We can derive the promised strengthening of Corollary 4.9. By the proof of [Bal09, Theorem 14.9] (originally [She99, II.1.6]):

Fact 4.15. *If \mathcal{K} is categorical in a $\lambda > H_2$, \mathcal{K} is (χ, H_2) -weakly tame for some $\chi < H_1$, and the model of size λ is χ -saturated, then \mathcal{K} is categorical in H_2 .*

Theorem 4.16. *If \mathcal{K} is categorical in a $\lambda \geq h(H_2^+)$, then \mathcal{K} is categorical in H_2 .*

Proof. By Theorem 4.14, there exists $\chi < H_1$ such that \mathcal{K} is (χ, H_2) -weakly tame. By Corollary 4.7, the model of size λ is χ -saturated. Now apply Fact 4.15. □

Remark 4.17. *We can replace H_2 above by any collection cardinal, see [Bal09, Definition 14.5] and the proof of Theorem 14.9 there.*

5. GOOD FRAMES AND WEAK TAMENESS

In [She09, Definition II.2.1]⁶, Shelah introduces *good frames*, a local notion of independence for AECs. This is the central concept of his book and has seen many other applications, such as a proof of Shelah's categoricity conjecture for universal classes [Vasc]. A *good μ -frame* is a triple $\mathfrak{s} = (\mathcal{K}_\mu, \perp, \text{ga-S}^{\text{bs}})$ where:

- (1) \mathcal{K} is a nonempty AEC which has μ -amalgamation, μ -joint embedding, no maximal models, and is stable in μ .
- (2) For each $M \in \mathcal{K}_\mu$, $\text{ga-S}^{\text{bs}}(M)$ (called the set of *basic types* over M) is a set of nonalgebraic Galois types over M satisfying (among others) the *density property*: if $M < N$ are in \mathcal{K}_μ , there exists $a \in |N| \setminus |M|$ such that $\text{ga-tp}(a/M; N) \in \text{ga-S}^{\text{bs}}(M)$.
- (3) \perp is an (abstract) independence relation on types of length one over models in \mathcal{K}_λ satisfying the basic properties of first-order forking in a superstable theory: invariance, monotonicity, extension, uniqueness, transitivity, local character, and symmetry (we will not give their exact meaning here).

As in [She09, Definition II.6.35], we say that a good μ -frame \mathfrak{s} is *type-full* if for each $M \in \mathcal{K}_\mu$, $\text{ga-S}^{\text{bs}}(M)$ consists of *all* the nonalgebraic types over M . We focus on type-full good frames in this paper. Given a type-full good μ -frame $\mathfrak{s} = (\mathcal{K}_\mu, \perp, \text{ga-S}^{\text{bs}})$ and $M_0 \leq M$ both in \mathcal{K}_μ , we say that a nonalgebraic type $p \in \text{ga-S}(M)$ *does not \mathfrak{s} -fork over M_0* if it does not fork over M_0 according to the abstract independence relation \perp of \mathfrak{s} . We say that a good μ -frame \mathfrak{s} is *on \mathcal{K}_μ* if its underlying class is \mathcal{K}_μ .

It was pointed out in [Vasa] (and further improvements in [Vasb, Section 10] or [VV, Theorem 6.12]) that tameness can be combined with superstability to build a good frame. At a meeting in the winter of 2015 in San Antonio, the first author asked whether weak tameness could be used instead. This is not a generalization for the sake of generalization because weak tameness (but not tameness) is known to follow from categoricity. As it turns out, the methods of [VV] can be used to answer in the affirmative:

Theorem 5.1. *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, \lambda]$ and has λ -symmetry.*

⁶We do *not* mention Shelah's axiom (B) requiring the existence of a superlimit model of size μ . In fact several papers (e.g. [JS13]) define good frames without this assumption. However the good frame that we build here is categorical in μ , hence has a superlimit.

If \mathcal{K} is (μ, λ) -weakly tame, then there exists a type-full good λ -frame with underlying class the saturated models in \mathcal{K}_λ .

Proof. First observe that limit models in \mathcal{K}_λ are unique (by Fact 2.7), hence saturated. By Fact 2.9, \mathcal{K} has χ -symmetry for every $\chi \in [\mu, \lambda]$. By Fact 2.8, for every $\chi \in [\mu, \lambda)$, $\mathcal{K}^{\chi^+ \text{-sat}}$, the class of χ^+ -saturated models in $\mathcal{K}_{\geq \chi^+}$ is an AEC with $\text{LS}(\mathcal{K}^{\chi^+ \text{-sat}}) = \chi^+$. Therefore (see [VV, Lemma 6.7]) $\mathcal{K}^{\lambda \text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\lambda \text{-sat}}) = \lambda$. By the λ -superstability assumption, $\mathcal{K}_\lambda^{\lambda \text{-sat}}$ is nonempty, has amalgamation, no maximal models, and joint embedding. It is also stable in λ . We want to define a type-full good λ -frame \mathfrak{s} on $\mathcal{K}_\lambda^{\lambda \text{-sat}}$. We define forking in the sense of \mathfrak{s} (\mathfrak{s} -forking) as follows: For $M \leq N$ saturated of size λ , a nonalgebraic $p \in \text{ga-S}(N)$ does not \mathfrak{s} -fork over M if and only if it does not μ -fork over M .

Now most of the axioms of good frames are verified in Section 4 of [Vasa], the only properties that remain to be checked are extension, uniqueness, and symmetry. Extension is by Fact 2.12.(3), and uniqueness is by uniqueness in μ (Fact 2.12.(1)) and the weak tameness assumption. As for symmetry, we know that λ -symmetry holds, hence we obtain the result by Section 3 of [VV]. \square

Remark 5.2. *If $\lambda = \mu^+$ above, then the hypotheses reduce to “ \mathcal{K} is superstable in μ and μ^+ and \mathcal{K} has μ^+ -symmetry”.*

Of course we can now combine this construction with our previous results:

Corollary 5.3. *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, h(\lambda))$. If \mathcal{K} is (μ, λ) -weakly tame, then there exists a type-full good λ -frame with underlying class the saturated models in \mathcal{K}_λ .*

Proof. Combine Theorem 5.1 and Theorem 3.10. \square

We obtain that a good frame can be built from categoricity in a high-enough cardinal (of arbitrary cofinality).

Corollary 5.4. *Let $\mu \geq H_1$. Assume that \mathcal{K} has no maximal models and is categorical in a $\lambda > \mu$. If the model of size λ is μ^+ -saturated (e.g. if $\text{cf}(\lambda) > \mu$ or by Corollary 4.7 if $\lambda \geq h(\mu^+)$), then there exists a type-full good μ -frame with underlying class the saturated models in \mathcal{K}_μ .*

Proof. By Fact 4.13, there exists $\chi < H_1$ such that \mathcal{K} is (χ, μ) -weakly tame. By the Shelah-Villaveces theorem (Fact 4.4), \mathcal{K} is superstable in every $\chi' \in [\chi, \lambda)$. By the proof of Fact 1.4 (which only uses that

the model of size λ is μ^+ -saturated), \mathcal{K} has μ -symmetry. Now apply Theorem 5.1 with (μ, λ) there standing for (χ, μ) -here. \square

REFERENCES

- [Bal09] John T. Baldwin, *Categoricity*, University Lecture Series, vol. 50, American Mathematical Society, 2009.
- [BG] Will Boney and Rami Grossberg, *Forking in short and tame AECs*, Preprint. URL: <http://arxiv.org/abs/1306.6562v9>.
- [BGKV] Will Boney, Rami Grossberg, Alexei Kolesnikov, and Sebastien Vasey, *Canonical forking in AECs*, Preprint. URL: <http://arxiv.org/abs/1404.1494v2>.
- [BK09] John T. Baldwin and Alexei Kolesnikov, *Categoricity, amalgamation, and tameness*, Israel Journal of Mathematics **170** (2009), 411–443.
- [Bon] Will Boney, *Computing the number of types of infinite length*, Notre Dame Journal of Formal Logic, To appear. URL: <http://arxiv.org/abs/1309.4485v2>.
- [Bon14] ———, *Tameness from large cardinal axioms*, The Journal of Symbolic Logic **79** (2014), no. 4, 1092–1119.
- [BV] Will Boney and Sebastien Vasey, *Chains of saturated models in AECs*, Preprint. URL: <http://arxiv.org/abs/1503.08781v3>.
- [GV] Rami Grossberg and Sebastien Vasey, *Superstability in abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1507.04223v2>.
- [GV06a] Rami Grossberg and Monica VanDieren, *Categoricity from one successor cardinal in tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), 181–201.
- [GV06b] ———, *Galois-stability for tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 1, 25–49.
- [GVV] Rami Grossberg, Monica VanDieren, and Andrés Villaveces, *Uniqueness of limit models in classes with amalgamation*, Mathematical Logic Quarterly, To appear. URL: <http://arxiv.org/abs/1507.02118v1>.
- [HS90] Bradd Hart and Saharon Shelah, *Categoricity over P for first order T or categoricity for $\phi \in L_{\omega_1, \omega}$ can stop at \aleph_k while holding for $\aleph_0, \dots, \aleph_{k-1}$* , Israel Journal of Mathematics **70** (1990), 219–235.
- [JS13] Adi Jarden and Saharon Shelah, *Non-forking frames in abstract elementary classes*, Annals of Pure and Applied Logic **164** (2013), 135–191.
- [She75] Saharon Shelah, *The lazy model theoretician’s guide to stability*, Logique et Analyse **18** (1975), 241–308.
- [She90] ———, *Classification theory and the number of non-isomorphic models*, 2nd ed., Studies in logic and the foundations of mathematics, vol. 92, North-Holland, 1990.
- [She99] ———, *Categoricity for abstract classes with amalgamation*, Annals of Pure and Applied Logic **98** (1999), no. 1, 261–294.
- [She09] ———, *Classification theory for abstract elementary classes*, Studies in Logic: Mathematical logic and foundations, vol. 18, College Publications, 2009.
- [SV99] Saharon Shelah and Andrés Villaveces, *Toward categoricity for classes with no maximal models*, Annals of Pure and Applied Logic **97** (1999), 1–25.

- [Vana] Monica VanDieren, *Superstability and symmetry*, Preprint. URL: <http://arxiv.org/abs/1507.01990v1>.
- [Vanb] ———, *Union of saturated models in superstable abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1507.01989v1>.
- [Van06] ———, *Categoricity in abstract elementary classes with no maximal models*, *Annals of Pure and Applied Logic* **141** (2006), 108–147.
- [Van13] ———, *Erratum to "Categoricity in abstract elementary classes with no maximal models" [Ann. Pure Appl. Logic 141 (2006) 108-147]*, *Annals of Pure and Applied Logic* **164** (2013), no. 2, 131–133.
- [Vasa] Sebastien Vasey, *Forking and superstability in tame AECs*, *The Journal of Symbolic Logic*, To appear. URL: <http://arxiv.org/abs/1405.7443v3>.
- [Vasb] ———, *Independence in abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1503.01366v4>.
- [Vasc] ———, *Shelah's eventual categoricity conjecture in universal classes*, Preprint. URL: <http://arxiv.org/abs/1506.07024v6>.
- [VV] Monica VanDieren and Sebastien Vasey, *Transferring symmetry downward and applications*, Preprint. URL: <http://arxiv.org/abs/1508.03252v2>.

DEPARTMENT OF MATHEMATICS, ROBERT MORRIS UNIVERSITY, MOON TOWNSHIP PA 15108

E-mail address: vandieren@rmu.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PENNSYLVANIA, USA

E-mail address: sebv@cmu.edu

URL: <http://math.cmu.edu/~svasey/>